

## References and studies

Hania B Bednarski, DO, FACOS

<https://link.springer.com/article/10.1007/s00520-018-4418-4> It still affects our economic situation: long-term economic burden of breast cancer and lymphedema; Lorraine T. Dean et al; Supportive Care in Cancer (2019) 27:1697–1708

<https://pubmed.ncbi.nlm.nih.gov/30521080/> : Early surveillance is associated with less incidence and severity of breast cancer-related lymphedema compared with a traditional referral model of care. Louise A Koelmeyer et al. Cancer. 2019 Mar 15;125(6):854-862.

<https://www.nature.com/articles/s41416-020-0844-4.pdf> : Increases in arm volume predict lymphoedema and quality of life deficits after axillary surgery: a prospective cohort study; Nigel Bundred et al; British Journal of Cancer

<https://pubmed.ncbi.nlm.nih.gov/32379872/> : Tissue Dielectric Constant Measures in Women With and Without Clinical Trunk Lymphedema Following Breast Cancer Surgery: A 78-Week Longitudinal Study; Linda A Koehler, Harvey N Mayrovitz; Phys Ther. 2020 Aug 12;100(8):1384-1392. doi: 10.1093/ptj/pzaa080.

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5737818/pdf/nihms925117.pdf> Breast Cancer Related Lymphedema risk is related to multidisciplinary treatment and not surgery alone – results from a large cohort study; Toan T Nguyen et al; Ann Surg Oncol. 2017 October ; 24(10): 2972–2980. doi:10.1245/s10434-017-5960-x

<https://pubmed.ncbi.nlm.nih.gov/25607768/>  
Immediate Implant Reconstruction Is Associated With a Reduced Risk of Lymphedema Compared to Mastectomy Alone: A Prospective Cohort Study; Cynthia L Miller et al. Ann Surg. 2016 Feb;263(2):399-405. doi: 10.1097/SLA.0000000000001128.

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5389857/>  
Sentinel Lymph Node Dissection With and Without Axillary Dissection in Women With Invasive Breast Cancer and Sentinel Node Metastasis: A Randomized Clinical Trial. Armando E Giuliano et al. JAMA. 2011 Feb 9; 305(6): 569–575. doi: 10.1001/jama.2011.90

<https://pubmed.ncbi.nlm.nih.gov/20863759/>  
Sentinel-lymph-node resection compared with conventional axillary-lymph-node dissection in clinically node-negative patients with breast cancer: overall survival findings from the NSABP B-32 randomised phase 3 trial. David N Crag et al. Lancet Oncol. 2010 Oct;11(10):927-33. doi: 10.1016/S1470-2045(10)70207-2.

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4354953/>

Axillary Reverse Mapping: Five Year Experience. Daniela Ochoa et al. *Surgery*. 2014 Nov; 156(5): 1261–1268. Published online 2014 Oct 17. doi: 10.1016/j.surg.2014.05.011

<https://pubmed.ncbi.nlm.nih.gov/18471751/>

Axillary reverse mapping: mapping and preserving arm lymphatics may be important in preventing lymphedema during sentinel lymph node biopsy. Cristiano Boneti et al. *J Am Coll Surg*. 2008 May;206(5):1038-42; discussion 1042-4. doi: 10.1016/j.jamcollsurg.2007.12.022. Epub 2008 Mar 3.

<https://clinicaltrials.gov/ct2/show/NCT03927027>

Axillary Reverse Mapping in Preventing Lymphedema in Patients With Breast Cancer Undergoing Axillary Lymph Node Dissection. ClinicalTrials.gov Identifier: NCT03927027

<https://clinicaltrials.gov/ct2/show/NCT03073096>

LYMPHA: Eliminating the Burden of Lymphedema in Patients Requiring Nodal Dissection. ClinicalTrials.gov Identifier: NCT03073096

<https://www.cancer.org/content/dam/cancer-org/research/cancer-facts-and-statistics/annual-cancer-facts-and-figures/2021/cancer-facts-and-figures-2021.pdf>

<https://pubmed.ncbi.nlm.nih.gov/32513642/>

Phase 1 Lymfactivin® Study: Short-term Safety of Combined Adenoviral VEGF-C and Lymph Node Transfer Treatment for Upper Extremity Lymphedema. Hartiala, Pauliina et al. *J Plast Reconstr Aesthet Surg*. 2020 Sep;73(9):1612-1621. doi: 10.1016/j.bjps.2020.05.009. Epub 2020 May 18.

[https://journals.lww.com/prsgo/Fulltext/2021/01000/Long\\_term\\_Results\\_of\\_Microvascular\\_Lymph\\_Node.36.aspx](https://journals.lww.com/prsgo/Fulltext/2021/01000/Long_term_Results_of_Microvascular_Lymph_Node.36.aspx)

Long-term Results of Microvascular Lymph Node Transfer: Correlation of Preoperative Factors and Operation Outcome. Rannikko, Eeva H et al. *Plastic and Reconstructive Surgery - Global Open*: January 2021 - Volume 9 - Issue 1 - p e3354  
doi: 10.1097/GOX.0000000000003354

<https://pubmed.ncbi.nlm.nih.gov/18942656/>

Treatment of upper limb lymphedema with combination of liposuction, myocutaneous flap transfer, and lymph-fascia grafting: a preliminary study. Qi, Fazhi et al. *Microsurgery*. 2009;29(1):29-34. doi: 10.1002/micr.20567.

Image credit: <https://en.wikipedia.org/wiki/Lymphedema>

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4127597/>

Breast cancer-related lymphedema: Symptoms, diagnosis, risk reduction, and management *World J Clin Oncol*. 2014 Aug 10; 5(3): 241–247.  
Published online 2014 Aug 10. doi: 10.5306/wjco.v5.i3.241