

A patient guide  
to understanding  
and managing  
chronic swelling.



**Tactile**  
MEDICAL®

# What is lymphedema?

Most of us know the circulatory system pumps blood from the heart to the rest of the body. But many people are not aware that the body has a second transport system — **the lymphatic system**. The lymphatic system consists of the tissues and organs that produce and store cells that fight disease and infection. They include the bone marrow, thymus and spleen, as well as the nodes and vessels that carry protein-rich fluid called lymph.

Lymphedema, and conditions causing chronic swelling, is a progressive condition in which excessive protein-rich fluids accumulate in the interstitium, the space between the cells. The condition occurs from impairment of the lymphatic system's ability to drain fluids and proteins from the spaces in between cells and transport this fluid, known as lymph, back to the circulatory system. A compromised or non-functioning lymphatic system places you at risk of medical complications including recurrent skin infections known as cellulitis.



## PATIENTS SUFFERING FROM LYMPHEDEMA

20 million+ Americans suffer from chronic swelling due to lymphedema or chronic venous insufficiency.<sup>1</sup>



Untreated lymphedema can make life miserable for those it affects. When patients receive a timely diagnosis and effective treatment, their health and quality of life improve.

**DR. THOMAS O'DONNELL**

BENJAMIN ANDREWS EMERITUS PROFESSOR AND CHAIR OF SURGERY TUFTS UNIVERSITY SCHOOL OF MEDICINE

# Types of lymphedema

---

## Primary lymphedema

Primary lymphedema, often called congenital lymphedema, may be caused by congenital (hereditary) malformations of the lymphatic system. Although it is largely attributed to genetic causes, lymphedema is also classified as primary when no known cause can be identified. Specific forms include:

- **Milroy disease** (*congenital lymphedema*). This disorder begins in infancy and causes lymph nodes to form abnormally.
- **Meige disease** (*lymphedema praecox*). This disorder often causes lymphedema around puberty or during pregnancy, though it can occur until age 35.
- **Late-onset lymphedema** (*lymphedema tarda*). Seen in patients with under developed lymphatics. Usually begins after age 35 as existing lymphatics lose the ability to compensate over time.

## Secondary lymphedema

Secondary lymphedema is an acquired condition caused by a physical disruption to the lymphatic system. Main causes include:<sup>2</sup>

- Chronic venous insufficiency
- Cancer treatments (*removal of lymph nodes and/or radiation therapy*), benign or malignant tumor growth
- Accident, trauma or surgery
- Infection
- Obesity
- Filariasis (*infiltration by insects or parasites*)

# Symptoms

---

- Unresolved or chronic swelling (*edema*)
- Swelling on the top of the foot
- Swollen and squared-off toes
- Heaviness or tightness in the affected limb
- Restricted range of motion
- Lymph fluid discharge from the skin (*lymphorrhea*)
- Skin changes such as:
  - “Orange peel” skin appearance
  - Thick, rigid skin (*hyperkeratosis and fibrosis*)
  - Bumpy wrinkling of the skin (*papillomatosis*)
  - Ulcerations of the skin
- “Ski jump” toenails (*hypoplastic and concave*)

## Stemmer’s sign: An indication of lymphedema in ten seconds or less

The Stemmer’s sign test results in either a positive or negative indicator for lymphedema. Simply pinch and lift a fold of skin at the base of your second toe or middle finger. If you can pinch and lift your skin, the Stemmer’s sign is negative. If you can’t, the sign is positive. False positives are rare, however, a negative does not always rule out lymphedema.

# Treatment

There is currently no cure for lymphedema. If untreated, it can worsen — leading to increased swelling and pain, progressive tissue hardening (*fibrosis*), and recurrent infections (*cellulitis*). Lymphedema may also reduce mobility, dexterity, and range of motion, limiting daily activities and significantly increasing healthcare costs. However, effective symptom management is possible, and proper diagnosis is critical to achieving effective treatment.

Management of lymphedema is usually achieved with a combination of therapist-directed in-clinic treatment, daily use of compression garments and at-home self-management with proven pneumatic compression devices from Tactile Medical. With effective treatment, you can manage your symptoms, minimize infections, improve your health and quality of life, and reduce overall healthcare costs.<sup>2,3,4</sup>

Ask your healthcare provider for more information about a pneumatic compression device (PCD).



**NIMBL™**  
Basic PCD for arms or legs.




**FLEXITOUCH® PLUS**  
Advanced PCD for arms, legs, chest, trunk, or head and neck.

# Meet Kylee.™

Stay supported with Kylee, an easy-to-use mobile app that helps you:

- **LEARN** about lymphedema.
- **TRACK** your symptoms and treatments.
- **SHARE** your progress with your doctor.



Tactile Medical is a leader in developing and marketing at-home therapies for people suffering from underserved, chronic conditions including lymphedema, lipedema, chronic venous insufficiency and chronic pulmonary disease by helping them live better and care for themselves at home.

Let's work together.



**Individual results may vary.**

**Indications/contraindications:** Indications, contraindications, warnings, and instructions for use can be found in the product labeling supplied with each device.

**Caution:** Federal (USA) law restricts these devices to sale by or on the order of a licensed healthcare practitioner.

**References:**

1. Dean S, Valenti E, Hock K, Leffler J, Compston A, Abraham W. The Clinical Characteristics of Lower Extremity Lymphedema in 440 Patients. *Jour of Vasc Surg: Venous & Lymph Disorders*. ePublished Feb 2020. Data on file at Tactile Medical.
2. Brayton KM, Hirsch AT, O'Brien PJ, et al. (2014). Lymphedema Prevalence and Treatment Benefits in Cancer: Impact of a Therapeutic Intervention on Health Outcomes and Costs. *PLoS ONE* 9(12): e114597. doi:10.1371/journal.pone.0114597
3. Ridner SH, McMahon E, Dietrich MS, et al. Home-Based Lymphedema Treatment in Patients with and Without Cancer-Related Lymphedema. *Oncology Nursing Forum*. July 2008; Vol. 35(4): 671–680.
4. Muluk SC, Hirsch AT, Tafe EC. Pneumatic Compression Device Treatment of Lower Extremity Lymphedema Elicits Improved Limb Volume and Patient-reported Outcomes. *EJVES*. 2013; Vol. 46(4): 480–487.

**Tactile Medical**

3701 Wayzata Blvd, Suite 300  
Minneapolis, MN 55416 USA

**Customer Care**

**Text or Call: 612.355.5100**

Toll Free Phone: 833.3TACTILE (833.382.2845)

Fax: 612.355.5101 / Toll Free Fax: 866.435.3949

Email: [customerservice@tactilemedical.com](mailto:customerservice@tactilemedical.com)

Hours: 7 a.m. to 5:30 p.m. CT, Monday–Friday

[tactilemedical.com](http://tactilemedical.com)

**Tactile**  
MEDICAL®

Tactile Medical and Flexitouch are registered trademarks and Nimbl and Kylee are trademarks of Tactile Medical.

©2025 Tactile Medical. All rights reserved. 702003-001 Rev F 12/2025